

Automated Patient Assessments After Outpatient Surgery Using an Interactive Voice Response System

Alan J. Forster, MD, MSc; Rinda LaBranche, RN; Robert McKim, MSc; John Wylam Faught, MD; Thomas E. Feasby, MD; Selikke Janes-Kelley, RN; Kaveh G. Shojania, MD; and Carl van Walraven, MD, MSc

Patients who undergo outpatient surgery do not stay in a hospital facility the evening after their procedure. With advances in minimally invasive surgery and anesthetic techniques, the volume of outpatient surgery is increasing dramatically and accounts for the majority of surgeries.^{1,2} Although outpatient surgeries are generally safe, risk is increased in certain circumstances.³⁻⁵ Outpatient surgery complications can be especially damaging because they occur at the patients' home outside of a monitored setting.

Fortunately, most complications occurring after outpatient surgery are self-limited. The most common complications include symptoms such as pain, nausea and vomiting, bleeding, and headache.⁶ Although these symptoms often are of minor clinical importance,^{4,6,7} it is important to monitor them for 3 reasons. First, symptoms can herald severe underlying problems such as wound infections or postoperative myocardial infarctions. Second, these symptoms can progress to severe outcomes such as shock, respiratory failure, or death, especially if there is a delay in treatment.^{8,9} Finally, symptoms can cause anxiety and impede a rapid return to normal function.

One common method of assessing complications after outpatient surgery is a nurse callback program. In such programs, a nurse calls patients on the day after surgery and will subsequently coordinate treatment if a patient is experiencing problems.¹⁰⁻¹² A limitation of such programs is their expense. To reduce costs, one could automate aspects of it using an interactive voice response system (IVRS),¹³ which is a technology allowing patients to interact with databases using a telephone.¹⁴⁻²¹ The IVRS could make an initial screening call to all patients to determine whether they were having a problem. Then, the IVRS could notify a nurse to follow up with the few patients who were screened "positive." This practice would obviate the need for a nurse to call every patient, significantly reducing the monitoring program's expense.

Two issues need to be considered by decision makers when contemplating whether to implement an IVRS-based patient monitoring system. First, will patients accept the technology? If patients cannot use the IVRS, then the technology is not capable of improving monitoring. Second, will the IVRS-based monitoring result in improved

identification and management of complications? If the system does not alert providers to patient problems in a timely manner, then the system is not useful.

Objective: To test the feasibility and utility of an interactive voice response system (IVRS) for monitoring patients after outpatient surgery.

Methods: We studied consecutive patients undergoing gynecologic day surgery. The IVRS called patients on the first postoperative day and asked them if they were experiencing new problems. Feasibility was assessed in terms of call responses and acceptance by patients. Utility was measured in terms of the ability of the IVRS to identify adverse events (AEs), defined as procedure-related symptoms requiring a physician or hospital visit. We contacted patients 30 days later to elicit their perceptions of the IVRS and determine AE status.

Results: Follow-up was complete for 249 of 270 enrolled patients (92%). The IVRS successfully contacted 130 patients (52%). Of the 22 patients (17%) who required a follow-up phone call, 9 had a new problem related to surgery, 7 had new or worsening symptoms, 6 wanted to speak with a nurse, and 1 had a medication-related problem. Patients remembering the automated call (n = 96) reported the system easy to use (82%) and comprehend (86%). Most patients (68%) preferred the IVRS to a personal follow-up call (probability greater than 50%, $P < .001$). AEs occurred in 40 patients (16%; 95% confidence interval = 12%, 21%). The IVRS did not identify any AEs because 90% of these occurred after the automated call.

Conclusion: An IVRS-based method of monitoring outpatient surgery patients is feasible. To improve utility, calls must occur later than first postoperative day.

(*Am J Manag Care.* 2008;14(7):429-436)

For author information and disclosures, see end of text.

In this issue
Take-away Points / p435
www.ajmc.com
Full text and PDF

We designed this study to assess these factors. We were primarily interested in understanding issues related to feasibility to determine whether further study of IVRS to manage patients was merited. We also were interested in understanding the potential utility of IVRS in terms of managing complications, particularly its ability to capture the timing and types of outcomes patients experienced. Because these data were not previously described, this study was necessary for the design of future IVRS-based interventions.

METHODS

Design, Setting, and Patients

We used a prospective cohort design to study women undergoing outpatient gynecologic surgery at the Royal Alexandra Hospital in Edmonton, Alberta, Canada. The Royal Alexandra Hospital is a teaching hospital within the Capital Health Region of Alberta. It has an active outpatient surgery program, conducting more than 14,000 procedures annually. Consecutive patients undergoing surgery between April 19, 2006, and June 9, 2006, were eligible. We excluded patients if they did not have a telephone, did not speak English, had cognitive impairment, or did not provide informed consent. We enrolled patients on the day of their surgery as they waited for their procedure. The study was approved by the joint University of Alberta/Capital Health Ethics Review Board.

Intervention

The study intervention is illustrated in **Figure 1**. After patient consent, we collected demographic and medical data. When patients were discharged from the outpatient surgery unit, the study nurse entered the patient's ID, phone number, and discharge date into the IVRS; this information was physically stored on a server in the hospital's data center. The nurse performed data entry using a Web-accessible personal computer in the outpatient surgery unit. The IVRS called patients on the day after their discharge from the day surgery unit. If there was no response to the call, the IVRS reattempted the call every 2 hours until a response occurred. If no response occurred before 9:00 PM that day, the system stopped calling the patient.

When a call was answered, the IVRS asked a series of questions that required a YES/NO answer. The IVRS used speech recognition software to update the database and prompt further actions. The system first asked: "Are you the patient discharged from the Royal Alexandra Hospital yesterday?" If the person responded NO, then the IVRS thanked the individual and disconnected. If the person responded YES, IVRS asked 4 questions and stored the responses:

- "Since your surgery, have you had any new or worsening symptoms?"
- "Since your surgery, have you had any problems related to your surgery?"
- "Since your surgery, have you had any problems related to your medications?"
- "Would you like to speak to a nurse?"

The IVRS transferred the patient to the provincial triage program (called HealthLink) after their first YES response to 1 of these 4 questions. HealthLink is a call center with nurses available to respond to patient queries at all times. Calls are answered by nurses, who are trained to triage patients according to the patient's problem and prespecified call algorithms. Call center nurses have specific algorithms to deal with postoperative problems.

The IVRS we used was CallAssure. CallAssure can be run on any modern personal computer equipped with a telephony card and at least 2 analog lines. In addition, CallAssure requires access to an e-mail server (for notification messages) and, optionally, a printer (for reports). CallAssure uses Nuance 8.5 as its speech recognition software. Nuance reports an overall accuracy of 97%, but this rate is likely much higher for responses to questions requiring YES/NO answers.²² We did not use routines for dealing with grammar mismatch errors to keep the system as simple as possible. Also, in previous testing, we discovered that patients found it difficult to revert to touch-tone inputs as many people use handheld devices. In our experience, these devices made it very difficult to use touch-tone input.

Follow-up and Study Outcomes

Thirty days after hospital discharge, we manually contacted all patients by telephone to administer a telephone survey. This survey was completed independent of any knowledge pertaining to the IVRS call. Patients were asked their recollection of the automated call, their opinion about some aspects of the IVRS (eg, perceived usefulness), whether they had any new or worsening health problems after their surgery, and whether they required any visits to providers since their surgery.

We supplemented the telephone survey with information from the health region's administrative data and the HealthLink call logs. We searched the respective data systems for visits to hospitals, results for any laboratory tests, or calls to HealthLink by the study patients within the 30 days after the surgery date. When such encounters existed, we obtained relevant information from the system, including the date, time, and reason for the hospital visit; the specimen type and date, as well as the results for all laboratory tests; and the call dialog, date, and time of all calls to HealthLink.

This study's main goal was to assess the IVRS feasibility, which was done by measuring the IVRS's call statistics and the survey results. A secondary objective was to determine the potential utility of using an IVRS for monitoring patients for adverse events. The relevant outcome for this aspect of the study was an adverse event, which we defined as any procedure-related symptom or poor health outcome requiring a visit to a physician or hospital.

Statistics and Data Management


We used SAS version 9.1 (SAS Institute, Inc, Cary, NC) for all data analysis. We report descriptive statistics using median and interquartile range (IQR) for continuous variables and frequency distributions for categorical variables. For outcomes, we present the proportion of patients with the outcome of interest with 95% confidence intervals (CIs), which we calculated using the Wilson score method.²³

We used a multivariable logistic regression model to measure the association between IVRS call response and the likelihood of experiencing an adverse event while controlling for patient and surgery factors. We first measured the association of experiencing an adverse event with patient and surgery factors at the univariate level by using the χ^2 test or the Wilcoxon rank sum test, respectively. We entered variables into our multivariable model if they were statistically associated with adverse event status at the $P < .20$ level.

Role of the Sponsors

The sponsors did not have any role in the design, conduct, analysis, or write-up of the study.

■ **Figure 1. Study Intervention**



IVRS indicates interactive voice response system; Pt, patient; RAH, Royal Alexandra Hospital.

^aAutomated calls were repeated until answered.


^bHealthLink is the teletriage program. Unfortunately, all patients connected to HealthLink hung up while in the call queue waiting to speak to a nurse. This is indicated in the figure by the gray line with the X.

RESULTS

Figure 2 summarizes the study flow and the main results for all patients. We enrolled 270 of 312 eligible patients during the study period. Thirty-four patients were not enrolled because they were missed by the research nurse, as she could not approach all patients. Eight patients refused participation in the study because they did not wish to take part in a research project. Of the 270 enrolled patients, 4 were not entered into the IVRS because of administrative errors and another 17 were lost to follow-up (6%). Our results are based on the 249 (92%) patients for whom we have 30-day follow-up data.

Table 1 describes the characteristics of patients in our cohort. The median age was 38 years (IQR = 31-49 years).

■ **Figure 2.** Study Flow and Results



IVRS indicates interactive voice response system; RN, registered nurse.

Most women had no comorbidities apart from the indication for surgery, virtually none had functional disabilities affecting their ability to perform activities of daily living, and few had surgeries in the preceding 12 months. The vast majority of patients were discharged on the day of surgery. The most common surgical procedures involved an abdominal approach (eg, laparoscopic surgery for endometriosis) or deep perineal approach (eg, dilation and curettage, cone biopsy).

We present call flow data in Figure 2. The IVRS made 318 calls to the 249 patients in the study. Most patients received 1 call; however, the system called back until the call was answered by the patient. Thus, many patients received more than 1 call, including 16 patients who were called 5 times. Of the telephone calls to the 249 patients, 163 were answered. Of these 163 calls, 130 were picked up by an individual who responded YES to the question regarding whether they were the correct patient. If patients responded YES to this ques-

tion, then they were very likely to answer all of the remaining questions with YES or NO responses. Of the 130 patients, 129 (99%) answered at least 1 other automated survey question and 104 (80%) answered all automated survey questions. Note that the IVRS connected patients to the HealthLink after a YES response. Therefore, if a patient responded YES to any question, then she did not have the opportunity to answer remaining questions. In total, 119 of 130 (92%) patients responded to all the questions they were intended to answer.

A minority of patients indicated via the IVRS that they were having a problem or that they wanted to speak to a nurse (Figure 2). Of the 130 patients who were contacted successfully, 9 patients stated they were having a problem related to their surgery, 7 reported having new or worsening symptoms, 6 reported a desire to speak with a nurse, and 1 reported having problems related to medications. None of these patients spoke to a HealthLink nurse. This is because after patients

Postoperative IVRS Calls

were connected to the HealthLink service, they hung up before speaking with a nurse. We surmise that this may have occurred because our call dialog did not explicitly mention they would be placed on hold. Therefore, no actions were taken as a result of the automated call.

At the time of our 30-day follow-up call, 40 of 249 women (16%; 95% CI = 12%, 21%) told us they had experienced an adverse event (Figure 2, Table 2). These adverse events included wound infections (n = 16 [6%]; 95% CI = 4%, 10%), excessive pain (n = 9 [4%]; 95% CI = 2%, 11%), bleeding (n = 6 [2%]; 95% CI = 1%, 5%), and miscellaneous other types of problems (n = 9 [4%]; 95% CI = 2%, 11%). These other adverse events included problems related to anesthesia (superficial phlebitis, pharyngotracheitis) and nausea and vomiting. A third of all adverse events required a visit to a hospital (including emergency department visits), whereas the remainder were managed in private doctors' offices. Three quarters of adverse events required some form of corrective action, including prescription of antibiotics (for wound infections) and increased analgesia medications (for pain). The median onset of these adverse events was 10 days after the surgery (IQR = 3-14 days). Only 4 adverse events (10%) occurred within 1 day of the surgery.

When we asked patients about their perceptions of the IVRS at the 30-day telephone follow-up, only 96 of the 130 patients recalled receiving an automated call from the IVRS. Of these patients, 83 (86%) reported they understood the system, 79 (82%) reported they found it easy to use, and 72 (75%) found the automated call useful. When asked if they would prefer a person to call them rather than an automated call, only 31 (32%) said they would.

We analyzed factors associated with adverse events (Table 3). Based on univariate assessments, the following factors were associated with adverse event risk at the $P < .05$ level: number of surgeries in preceding year and the IVRS correctly contacting the patient. In a multivariable logistic regression model adjusting for age, the presence of diabetes mellitus, the number of surgeries in the preceding year, and whether the patient was released from the hospital on the day of surgery, we found that receiving a call from the IVRS was associated with a significant reduction in the probability of adverse event occurrence (odds ratio = 0.40; 95% CI = 0.19, 0.83). No other factor was independently associated with adverse event risk, including the actual responses provided to the IVRS.

DISCUSSION

We designed this study to determine the feasibility and utility of an IVRS to monitor patients following outpatient surgery. Overall, our data show that the system was feasible,

■ **Table 1. Patient Characteristics**

Characteristic	Value ^a
No. of patients	249
Age, median (interquartile range), y	38 (31-49)
Comorbidities	
Pulmonary disorders	23 (9)
Diabetes mellitus	8 (3)
Cardiac disease	9 (4)
No. of disabilities	
0	239 (96)
1	8 (3)
2+	2 (1)
No. of surgeries in preceding year	
0	207 (83)
1	37 (15)
2+	5 (2)
Surgical approach	
Superficial perineal	10 (4)
Deep perineal	117 (47)
Abdominal	122 (49)
Released from hospital on day of surgery	244 (98)
Lived in local region	199 (80)

^aValues are number (%) except where indicated.

with 52% of patients being reached by the IVRS. Almost all of these patients understood the call dialog and a majority preferred an automated call to a personal one.

Despite our optimism regarding feasibility, our results raised concerns related to the IVRS's utility. The system failed to connect patients to the teletriage program. This was because patients hung up while they waited to speak with a HealthLink nurse. This problem may be obviated by changing the IVRS call dialog to explicitly instruct patients to hold while a nurse is made available. We also found that patients experienced most of their adverse events after the IVRS call. This occurred because we designed our IVRS to call on the first postoperative day (ie, our design reflected existing work flows in most day surgery units). This problem is easily remedied by delaying the call or repeating it several days later. The most important finding of our study was that 1 in 6 women experienced an adverse event. This finding emphasizes the need for improved monitoring of patients undergoing outpatient surgery. Given the expense of having nurses perform monitoring, we believe our study supports a need for further efforts to adapt IVRS technologies for this task.

■ **Table 2.** Adverse Events Captured at 30-day Telephone Interview

Type of Adverse Event ^a	No. (%) (n = 249)	No. of Days Until Adverse Event, Median (IQR)	No. (%) Who Required Corrective Action
All types	40 (16)	10 (3-14)	30 (75)
Visit to a hospital required	13 (5)	4 (2-11)	9 (69)
Wound infection	16 (6)	10 (2-13)	15 (94)
Visit to a hospital required	2 (1)	6.5 (2-11)	2 (100)
Pain	9 (4)	10 (1-17)	6 (67)
Visit to a hospital required	5 (2)	3 (1-14)	3 (60)
Bleeding	6 (2)	13 (10-21)	3 (50)
Visit to a hospital required	3 (1)	10 (3-13)	2 (67)
Other	9 (4)	6.5 (3.5-9)	6 (67)
Visit to a hospital required	3 (1)	4 (0-6)	2 (67)

IQR indicates interquartile range.
^aAdverse events were procedure-related symptoms necessitating a physician or hospital visit.

the current business practice of manually calling patients does not require the patient to verify who they are. Second, we felt any technical solution would be associated with increasing complexity, which might decrease usability and patient acceptance.

We performed a well-designed study. Its strengths included very broad inclusion criteria, a small proportion of patients lost to follow-up, and clinically important outcomes captured in all patients. Our conclusions were weakened because we are unaware why some patients did not respond to the IVRS. Some potential reasons include technical factors (eg, the IVRS did not call

A follow-up telephone call to assess patients following outpatient surgery appears to be a good idea. However, there are limited data describing its rationale and benefit.^{11,24} Despite this, many health facilities and some health systems have a policy to attempt to call patients for this purpose. It is unknown how many facilities routinely contact patients after outpatient surgery, how effectively surgery units contact patients, and what follow-up occurs once a problem is identified. It is likely that facilities do not strictly adhere to the policy and do not track statistics for 2 reasons: a shortage of nursing staff to place the phone calls, and a lack of data systems to help support the work flow. The IVRS we tested could solve these 2 problems, while still being acceptable to patients.

Our study does not support the conclusion that IVRS calls improve patient outcomes after day surgery. The association of successful call completion with a lower adverse event risk is intriguing, but is almost certainly due to confounding. Although it is possible that patients may not have answered the telephone because they were having symptoms, we think this is unlikely given the adverse event timing we observed. It is remotely possible that patients who answered the IVRS were more compliant with follow-up instructions, which led to a reduced risk. This seems highly unlikely.

When we designed the IVRS, we debated whether methods to verify the true identity of a patient should be included. These methods could have required the patient to provide their birth date, the citation of a unique personal identification number at the time of enrollment, or the use of biometrics such as voice recognition. In our final design, we favored omitting these methods for 2 reasons. Most importantly,

patients as programmed) or patient factors (eg, patients may not have understood the call prompts). We think it is more likely that patients did not answer the phone when called, as this factor has explained previous unsuccessful attempts to contact patients using manual calls.^{24,25} Similarly, we do not know whether the people who answered the IVRS responded to questions truthfully. An indication of this possibility is that 34/130 (26%) patients could not recall the IVRS call when we personally interviewed them. However, we think it is unlikely that people gave untruthful answers because prior IVRS research evaluating correspondence between personal and IVRS interviews shows that patients generally give consistent answers using both approaches. It is very possible that given patients' busy lives, they might not recall a brief phone call (median duration = 60 s; IQR = 55-68 s) 4 or 5 weeks earlier. Regardless, even if all the patients who did respond to the IVRS but could not remember the call had negative opinions regarding the automated calls, the majority of patients in our study would still have favorable opinions of the technology.

Based on this study, we recommend several changes to the design of the telephone-based interventions to monitor patient safety following outpatient surgery (whether automated or not). First, patients should be called later than 1 day after the surgery. Ninety percent of adverse events started after the first postoperative day. Either a second call should be made several days later, or the first call needs to be made at that time. Second, more attempts to call patients are required. Many patients in our study did not respond despite several attempts by the system. Third, an enhanced response to patients requiring follow-up is essential. If one is going to automatically link

patients to a triage program, then the technology needs to work. In our study, we found that patients hung up after they were connected to the triage program, probably because they did not understand they had to wait to speak to a nurse. This problem could be fixed by explaining with a call script the need to hold for a nurse. Fourth, the outpatient surgery unit needs to be notified of the potential problem. It would not be too onerous for the unit to personally call these patients, as only 17% of patients said YES to the IVRS-based question (Figure 2). Alternatively, the unit could focus on the patients not responding to the IVRS, as these were the ones who appeared to be at higher risk of problems. Finally, patients should be able to make in-bound calls to the IVRS.

In conclusion, our data show that it is feasible to monitor patients following outpatient surgery using an IVRS. If a particular outpatient surgery unit requires proof that follow-up calls improve patient outcomes, then we recommend they wait for randomized trial evidence before implementation. However, if an outpatient unit feels that follow-up calls must be done to meet regulatory requirements or because they simply want to enhance contact with patients following day surgery, then, given our findings, using an IVRS-based approach appears justified, assuming minor modifications in the technology's design are made.

Table 3. Association of Patient and Call Characteristics With Patient Outcomes

Characteristic	Patient Experienced Adverse Event ^a		P ^b	OR ^c (95% CI)
	No	Yes		
No. of patients	209	40		
Age, median (IQR), y	39 (31-49)	35.5 (30-40)	.10	0.97 (0.94, 1.00)
Comorbidities				
Pulmonary disorders	19 (9)	4 (10)	.77	
Diabetes mellitus	5 (2)	3 (8)	.12	3.7 (0.79, 17.4)
Cardiac disease	9 (4)	0 (0)	.36	
No. of disabilities				
0	201 (96)	38 (95)	.67	
1+	8 (4)	2 (5)		
No. of surgeries in preceding year				
0	178 (85)	29 (73)	.05	2.2 (0.99, 5.1)
1+	31 (14)	11 (28)		
Surgical approach				
Abdominal	110 (53)	17 (43)	.24	
Other	99 (47)	23 (58)		
Released from hospital on day of surgery	206 (99)	38 (95)	.18	0.30 (0.04, 2.0)
Lived in local region	169 (81)	30 (75)	.4	
Correct patient called by IVRS system	116 (56)	14 (35)	.02	0.40 (0.19, 0.83)

CI indicates confidence interval; IQR, interquartile range; IVRS, interactive voice response system; OR, odds ratio.
^aValues are number (%) except where indicated.
^bUnivariate assessment of the probability of falsely concluding there is an association between adverse event risk and a specific patient characteristic.
^cMultivariate OR measuring the increased odds of experiencing an adverse event given the presence of a factor.

Take-away Points

To monitor for adverse events in patients undergoing outpatient surgery, many institutions have a policy for nurses to call patients on the first postoperative day, a process that is expensive and difficult to audit.

- An interactive voice response system (IVRS) can be used for postoperative calls.
- Use of the IVRS is more efficient and does not compromise the quality of the assessments.
- However, to improve utility, calls must occur later than the first postoperative day.

Author Affiliations: From the Department of Medicine, University of Ottawa (AJF, KGS, CVW), Ottawa, Ontario; the Clinical Epidemiology Program, Ottawa Health Research Institute (AJF, KGS, CVW), Ottawa, Ontario; Capital Health (RL, RM, WF, TEF, SJ-K), Edmonton, Alberta; the Research, Evaluation and Quality Improvement Center, University of Alberta (RM, SJ-K), Edmonton, Alberta; the Department of Obstetrics and Gynaecology, The Ottawa Hospital (WF), Ottawa, Ontario; and the Faculty of Medicine, University of Calgary (TEF), Calgary, Alberta.

Funding Source: This research was funded by the Canadian Patient Safety Institute, the Ottawa Hospital Center for Patient Safety, and Capital Health, Edmonton area. Vocantas Inc provided technical support for the project.

Author Disclosures: Dr Forster is supported by an Ontario Ministry of Health Career Scientist Award and is a coinventor on a patent for an interactive voice response system. Dr Shojania is supported by a Government of Canada Research Chair in Patient Safety and Quality Improvement. The other authors (RL, RM, JWF, TEF, SJ-K, CVW) report no relationship or

financial interest with any entity that would pose a conflict of interest with the subject matter of this article.

Authorship Information: Concept and design (AJF, RL, RM, JWF); acquisition of data (AJF, RL, RM, JWF, SJ-K); analysis and interpretation of data (AJF, TEF, KGS); drafting of the manuscript (AJF, KGS); critical revision of the manuscript for important intellectual content (AJF, RL, RM, JWF, TEF, KGS, CVW); statistical analysis (AJF, CVW); provision of study materials or patients (AJF, RL, RM, JWF, SJ-K); obtaining funding (AJF, RL, RM); administrative, technical, or logistic support (RL, TEF, SJ-K, CVW); and supervision (AJF, JWF).

Address correspondence to: Alan J. Forster, MD, MSc, Clinical Epidemiology Program, Ottawa Health Research Institute, 725 Parkdale Ave, Ottawa, Ontario K1Y 4E9, Canada. E-mail: aforster@ohri.ca.

REFERENCES

1. **Eggertson L.** Ten year trend: surgeries up, hospitalizations down. *CMAJ*. 2007;176(6):756.
2. **Aylin P, Williams S, Jarman B, Bottle A.** Trends in day surgery rates. *BMJ*. 2005;331(7520):803.
3. **Tham C, Koh KF.** Unanticipated admission after day surgery. *Singapore Med J*. 2002;43(10):522-526.
4. **Shnaider I, Chung F.** Outcomes in day surgery. *Curr Opin Anaesthesiol*. 2006;19(6):622-629.
5. **Twersky R, Fishman D, Homel P.** What happens after discharge? Return hospital visits after ambulatory surgery. *Anesth Anal*. 1997;84(2):319-324.
6. **Wu CL, Berenholtz SM, Pronovost PJ, Fleisher LA.** Systematic review and analysis of postdischarge symptoms after outpatient surgery. *Anesthesiology*. 2002;96(4):994-1003.
7. **Rawal N.** Postoperative pain treatment for ambulatory surgery. *Best Pract Res Clin Anaesthesiol*. 2007;21(1):129-148.
8. **Coldiron B, Fisher AH, Adelman E, et al.** Adverse event reporting: lessons learned from 4 years of Florida office data. *Dermatol Surg*. 2005;31(9, pt 1):1079-1092; discussion 1093.
9. **Grogaard B, Kimsas E, Raeder J.** Wound infection in day-surgery. *Ambul Surg*. 2001;9(2):109-112.
10. **Gilmartin J, Wright K.** The nurse's role in day surgery: a literature review. *Int Nurs Rev*. 2007;54(2):183-190.
11. **Fallis WM, Scurrah D.** Outpatient laparoscopic cholecystectomy: home visit versus telephone follow-up. *Can J Surg*. 2001;44(1):39-44.
12. **Burden N.** Telephone follow-up of ambulatory surgery patients following discharge is a nursing responsibility. *J Post Anesth Nurs*. 1992;7(4):256-261.
13. **Forster AJ, van Walraven C.** Using an interactive voice response system to improve patient safety following hospital discharge. *J Eval Clin Pract*. 2007;13(3):346-351.
14. **Biem HJ, Turnell RW, D'Arcy C.** Computer telephony: automated calls for medical care. *Clin Invest Med*. 2003;26(5):259-268.
15. **DeMolles DA, Sparrow D, Gottlieb DJ, Friedman R.** A pilot trial of a telecommunications system in sleep apnea management. *Med Care*. 2004;42(8):764-769.
16. **LeBaron CW, Starnes DM, Rask KJ.** The impact of reminder-recall interventions on low vaccination coverage in an inner-city population. *Arch Pediatr Adolesc Med*. 2004;158(3):255-261.
17. **Mahoney D, Tennstedt S, Friedman R, Heeren T.** An automated telephone system for monitoring the functional status of community-residing elders. *Gerontologist*. 1999;39(2):229-234.
18. **Stuart GW, Laraia MT, Ornstein SM, Nietert PJ.** An interactive voice response system to enhance antidepressant medication compliance. *Top Health Inf Manage*. 2003;24(1):15-20.
19. **Pinto BM, Friedman R, Marcus BH, Kelley H, Tennstedt S, Gillman MW.** Effects of a computer-based, telephone-counseling system on physical activity. *Am J Prev Med*. 2002;23(2):113-120.
20. **Leirer VO, Morrow DG, Pariente G, Doksum T.** Increasing influenza vaccination adherence through voice mail. *J Am Geriatr Soc*. 1989;37(12):1147-1150.
21. **Leirer VO, Morrow DG, Tanke ED, Pariente GM.** Elders' nonadherence: its assessment and medication reminding by voice mail. *Gerontologist*. 1991;31(4):514-520.
22. **Nuance.** Nuance 8.5 specifications. <http://www.nuance.com/recognizer/nuancerecognizer>. Accessed April 22, 2008.
23. **Newcombe RG.** Two-sided confidence intervals for the single proportion: comparison of seven methods. *Stat Med*. 1998;17(8):857-872.
24. **Dewar A, Scott J, Muir J.** Telephone follow-up for day surgery patients: patient perceptions and nurses' experiences. *J Perianesth Nurs*. 2004;19(4):234-241.
25. **Dudas V, Bookwalter T, Kerr KM, Pantilat SZ.** The impact of follow-up telephone calls to patients after hospitalization. *Dis Mon*. 2002;48(4):239-248. ■